

Syphilis Morbidity in Yakima County

2017-2021 YTD

	2017	2018	2019	2020	2021 YTD
Overall Morbidity					
Total syphilis (all stages)*	36	50	41	50	163
Primary & Secondary	14	12	8	22	79
Early Latent (non-primary non-secondary)	10	23	8	10	40
Late or unknown duration	12	15	25	18	35
Congenital	0	0	0	0	9
Gender†					
Male	25	37	29	27	95
Female	11	13	11	23	67
Race/Ethnicity					
White non-Hispanic	+	15	9	5	37
Black non-Hispanic	0	0	+	+	+
Asian, non-Hispanic	+	+	0	0	0
AIAN, non-Hispanic	0	+	0	+	21
NHOPI, non-Hispanic	0	0	0	0	0
Multi, non-Hispanic	+	+	+	+	+
Hispanic	24	27	29	28	78
Unknown	+	+	+	+	14
Risk Factors					
MSM	16	23	14	11	28
WSM	11	13	11	23	58
Pregnant	+	+	+	+	18
PLWH	+	+	+	+	13
Drug Use‡	+	+	+	+	29

Counts less than 10, or where small numbers can be deduced using subtraction from other cells, have been suppressed to protect data confidentiality.

2021 YTD preliminary data includes cases reported between Jan 01, 2021 and Dec 31, 2021. This data is subject to change.

**All stages of syphilis includes primary, secondary, early latent (non-primary non-secondary), late/unknown duration, and congenital*

†For the purposes of this analyses, only cis-gender males and females were included

‡Drug use includes any reported use of one or more of the following: heroin, meth, crack, IDU, cocaine, nitrates, hallucinogens, alcohol, opiates, GHB, ketamine, ecstasy, marijuana

Source: PHIMS-STD 06/01/2021